

Khoo, Yvonne J-Lyn (2008) CYP2C8 Polymorphisms among malaria patients in Guinea-Bissau. In: 2nd Health and Medical Sciences Conference / 2nd Penang International Postgraduate Convention: Chemical Biology, Innovative Exploration of New Horizons in Science, 18-20 June 2008, University of Science, Malaysia. (Unpublished)

Downloaded from: <http://insight.cumbria.ac.uk/id/eprint/4994/>

Usage of any items from the University of Cumbria's institutional repository 'Insight' must conform to the following fair usage guidelines.

Any item and its associated metadata held in the University of Cumbria's institutional repository Insight (unless stated otherwise on the metadata record) may be copied, displayed or performed, and stored in line with the JISC fair dealing guidelines (available [here](#)) for educational and not-for-profit activities

provided that

- the authors, title and full bibliographic details of the item are cited clearly when any part of the work is referred to verbally or in the written form
 - a hyperlink/URL to the original Insight record of that item is included in any citations of the work
- the content is not changed in any way
- all files required for usage of the item are kept together with the main item file.

You may not

- sell any part of an item
- refer to any part of an item without citation
- amend any item or contextualise it in a way that will impugn the creator's reputation
- remove or alter the copyright statement on an item.

The full policy can be found [here](#).

Alternatively contact the University of Cumbria Repository Editor by emailing insight@cumbria.ac.uk.



Erasmus Mundus



CYP2C8 Polymorphisms among malaria patients from Guinea-Bissau

Yvonne Khoo J-Lyn

Supervised by Prof. Dr. Vera Ribeiro

European Joint Master in Water and Coastal Management

Universidade do Algarve

Portugal

19 June 2008




CONTENTS

- Introduction
- Research objectives
- Materials and methods
- Results
- Discussion
- Conclusion
- Future recommendations
- Bibliography
- Acknowledgements

INTRODUCTION



Malaria is one of the major public health problems in more than 90 countries, inhabited by a total of some 2.4 billion people, representing about 40% of the world's population (WHO, 2004).





Malaria endemic areas

■ Distribution of Malaria




CDC

Amodiaquine (AQ) has been recently introduced into artemisinin-based combination therapy for use in malaria control programmes and as a first line treatment for children with uncomplicated malaria (WHO, 2006).




Besides amodiaquine, *CYP2C8* also metabolizes several therapeutically important drugs and endogenous substances including..




- paclitaxel
- verapamil
- rosiglitazone
- cerivastatin
- amiodarone
- dapsone
- all-*trans*-retinoic acid
- arachidonic acid





CYP2C8 is mainly expressed in the liver, as well as in various extrahepatic tissues such as the vascular smooth muscles (Klose et al., 1999; Fleming, 2001).

The main *CYP2C8* polymorphisms known code for the amino acid changes I269F, R139K, K399R and I264M.



These SNPs define 3 main non-wild-type alleles: *CYP2C8****2**, *CYP2C8****3** and *CYP2C8****4**.

A glance at Guinea-Bissau



Source: travelpod.com

A glance at Guinea-Bissau



Source: travelpod.com



Canchungo hospital, Guinea-Bissau

Source: www.kalpana.it

RESEARCH OBJECTIVES



- To **study *CYP2C8* alleles** among malaria patients from Guinea Bissau
- To **assist policy-makers** in the management of malaria in Guinea-Bissau
- To **generate pharmacogenetic data** for the evaluation of treatment and drug dispensation
- To **contribute findings** to other databases and bio-banks within and outside Europe
- To allow **further comparisons** with other populations previously characterized in the Center for Molecular and Structural Biomedicine, Universidade do Algarve, Portugal

MATERIALS AND METHODS



Subjects : 91 randomly selected malaria patients from Guinea-Bissau



DNA Extraction



Polymerase Chain Reaction (PCR)



Restricted Fragment Length Polymorphism (RFLP)




Statistical Analysis



Allelic frequencies determined - PM alleles? EM alleles? etc

RESULTS



Lane 1: ϕ X174 DNA/HinfI Marker; Lane 2: Homozygous mutant for the *CYP2C8**2 allele; Lane 3, 5, 6: Homozygous wild-type for the *CYP2C8**2 allele; Lane 4: Heterozygous for for *CYP2C8**2 allele; Lanes 7 to 11: Homozygous wild-type bands for the *CYP2C8**4 variant; Lane 12: PCR amplicon used to generate the RFLPs


***CYP2C8* allele frequencies obtained:**

$$CYP2C8^*2 = 0.2418$$

$$CYP2C8^*3 = 0.3242$$

$$CYP2C8^*4 = \text{not detected}$$

Distribution of *CYP2C8* genotypes among GB subjects



DISCUSSION



- Comparison of ***CYP2C8* genotypes** with other populations
- Comparison of ***CYP2C8* allele frequencies** with other populations
- Comparison of ***CYP2C8* allele frequencies** between malaria patients from **GB** and **Zanzibar**

Comparison of *CYP2C8* allele frequencies between malaria patients from Guinea-Bissau and Zanzibar




- Higher prevalence of the *CYP2C8**3 allele in West Africa

Comparison with Asian and Oceanic *CYP2C8* allele frequencies




- Significant differences not detected

CYP2C8 genotype comparison with 5 other populations




Comparison of *CYP2C8* allele frequencies with other populations



CONCLUSION



- Highest frequency of *CYP2C8* variant alleles ever recorded in a population of African descent.
- High occurrence of *CYP2C8**2 and *CYP2C8**3 alleles among malaria patients in Guinea-Bissau.
- This implies a high incidence of *CYP2C8* poor metabolizer alleles among malaria patients in Guinea-Bissau who may be at a greater risk of adverse effects compared to other populations previously characterized.



FUTURE RECOMMENDATIONS



- Further investigation taking into account the effects of *CYP2C8* metabolism on the pharmacokinetics of antimalarials



- Study of polymorphisms in healthy subjects

BIBLIOGRAPHY

Bradley, D.J., 1974. Chapter in Human Rights in Health. Ciba Foundation Symposium **23**: 81–98.

Cavaco, I., Stromberg-Norklit, J., Kaneko, A., Msellem, M.I., Dahoma, M., Ribeiro, V.L., Bjorkman, A. and Gil, J.P., 2005. *CYP2C8* polymorphism frequencies among malaria patients in Zanzibar. *Eur J Clin Pharmacol* **61**: 15–18

Cavaco, I., Piedade, R., Gil, J.P. and Ribeiro, V., 2006. *CYP2C8* polymorphism among the Portuguese. *Clin Chem Lab Med* **44**: 168–170.

Kofoed, P.E., 2006. Treatment of uncomplicated malaria in Guinea-Bissau. PhD Thesis. Karolinska Institutet, Stockholm, Sweden.

WHO, 2006. WHO Guidelines for the treatment of malaria. WHO Press, Geneva.



Special thanks to my supervisor
Prof. Dr. Vera Ribeiro

My Erasmus coordinator Prof. Dr
Alice Newton

Prof. Virgílio do Rosário

This research involves a
collaboration with the Center of
Malaria and Tropical Diseases,
Universidade de Lisboa, in the
frame of the Portuguese Network for
Malaria and the Network for Tropical
Health Research in Portuguese-
speaking countries (RIDES-PLP).

This project is partially supported by
the Project Ceratonia from Caixa
Geral de Depósitos.

Khoo YJL is a recipient of an
Erasmus Mundus studentship.

Muito obrigada